

IDRC-LIN TaaTT OPPERATE

-

.

INTERNATIONAL WORKSHOP ON MOUNTAIN CROPS

AND GENETIC RESOURCES

KATHMANDU, NEPAL

FEBRUARY 16-19, 1987

THE ANDEAN PHYTOGENETIC AND ZOOGENETIC RESOURCES

Mario E. Tapia, Agr. Eng. PhD, INIPA/IDRC/ACDI Lima, Perú

Nicolás Mateo, Agr. Eng. PhD, IDRC Canada

ARCHIU 11 FTE) 200 10

October, 1986

TOPIC II

THE ANDEAN PHYTOGENETIC AND ZOOGENETIC RESOURCES

CONTENTS

- I. THE ANDES AS CENTER OF ORIGIN OF CROP AND ANIMAL SPECIES
- II. THE ANDEAN MOUNTAINS AND THEIR GENETIC DIVERSITY

CROPS

- Grains
- Tubers
- Roots

ANIMALS

- South American Camelids

and water and the second se

- Guinea pigs ("cuys")
- III. COLLECTION EXPEDITIONS : THE ANDEAN NATIVE CROPS AND DOMESTIC ANIMAL RESOURCES GERMPLASM BANKS.
- IV. PRESENT AND POTENTIAL USE OF PHYTOGENETIC AND ZOOGENETIC RESOURCES

THE ANDEAN PHYTOGENETIC AND ZOOGENETIC RESOURCES

"A country is a given technological capacity over its geography and it is only through accepting and being aware of such geography that we will make every effort to produce the food that such geography may offer us. This is the only means to break the vicious circle of imports which diminishes the wealth of our peasants and limits urban industry to insuficient markets".

Mr. Alan García Pérez President of the Republic of Perú

The Andean mountains cross from North to South of the South American continent and embrace tropical and subtropical latitudes having altitudes which range from sea level to over 4,000 m. The area located above 2,000 m occupies more than 2'000,000 hectares belonging to seven different countries.

Within the framework of the above conditions, settlers who arrived 10,000 years ago used this broad range of flora to select species such as : grains, tubers, roots and nourishing cormos, spice and healing plants, and fibers for their clothing.

The ample range of pasture in the highlands, allowed for the domestication of wild ruminants, as the New World camelids (llama and alpaca), whose meat, skin and fibers, were used. Likewise, old chronicles do mention the . domestication of a type of hen and bird such as the "huallata" or South American goose.

I. THE ANDES AS CENTER OF ORIGIN OF CROP AND ANIMAL SPECIES

Vavilov (1935) refers to the Andean region of Ecuador, Perú and Bolivia, as one of the eight centers of origin of crop plants in the world. This region was the center of the Pre-Incan civilizations and gave birth to the Incan Empire in the XI century. In spite of the fact that this zone is not significantly large, when compared to South America as a whole, it is of utmost importance for endemic crop plants and animal resources.

Yacovleff and Herrera (1943) reviewed and analyzed the information compiled by Spanish chronicle authors since their arrival to the Andes, on plants demosticated by natives of the pre-Spanish culture. Mention is made of 159 species being used, even though some of them are found in semi-wild state.

Table No. 1 shows the main domesticated crop species which could have potential use in other regions of the world.

The domestication of wild animal species in the Andes was focused on animals which could provide both their skin and fiber and be used as a means of transportation for food and other materials. The South American camelids include two domestic species and two wild species adapted to the different conditions of the pastures of the Andes.

TABLE 1 -

MAIN ANDEAN NATIVE FOOD SPECIES ABOVE 2,000 M *

COMMON NAME	SCIENTIFIC NAME	BUTANICAL FAMILY
GRAINS Quinoa Kañīwa Kiwîcha Tarwî	Chenopodium quinoa Chenopodium pallidicaule Amaranthus caudatus Lupinus mutabilis	Quenopodiaceae Quenopodiaceae Amarantaceae Leguminosae
TUBERS Oca Olluco Mashua	Oxalis tuberosa Ullucus tuberosus Tropaeolum tuberosum	Oxalidaceae Baselaceae Tropaeolaceae
ROOTS Maca Jimaca, Ajipa Arracacha Yacon "Jiguima"	Lepidium meyenii Pachurrhizus ahipa Arracacia xanthorrhiza Polymnia sonchifolia	Cruciferae Leguminosae Umbeliferae Compositae
COMMON FRUITS Nuez del Perú Chirimoya Pacaæ, "guamos" Huagra-manzana Capulí Mora de Castilla	Juglans peruyiana Annona cherimola Inga feuillei Crataegus stipulosa Prunus serotina Rubus glaucus	Juglandaceae Anonaceae Leguminosae Rosaceae Rosaceae Rosaceae
OTHER FRUITS Ciruela del Fraile Tumbo, curuba Tíntin Lucuma Capulí, "uchuba" Sacha tomate	Bunchosia armeniaca Passiflora mollisima Passiflora pinnatistipula Lucuma bifera Pnysalis peruviana Cyphomandra betacea	Malpigiaceae Pasifloraceae Pasifloraceae Sapotaceae Solanaceae Solanaceae

*Maize and potato are not mentioned due to their worldwide distribution.

٠

~

-

•

TABLE 2

MAIN ANDEAN ANIMAL SPECIES

SPECIES	SCIENTIFIC NAME	PRESENT STATUS
MAMMALS		
Llama Alpaca Vicuña Huanaco	Lama glama Lama glama pacos Lama glama vicugna Lama glama guanicol	Domesticated Domesticated Wild Wild
CERVIDAE		
Déer	Hippocamelus antisensis	Wild
CARNIVORS		,
Fox Puma	Dusicyon sp Felis concolor	Wild Wild
POULTRY		• .
Condor Wallata Pariwana Partridge	Vulture gryphus Choephaga melanoptera Phoenicoparus andinas Nothoprocta sp	Wild Wild Aild Wild
RODENTS		•
Cuy Vizcacha Chinchilla	Cavia parcellus Lagidium peruanum Chinchilla Taniger	Domesticated Wild Domesticated

II. THE ANDEAN MOUNTAINS AND THEIR GENETIC DIVERSITY

CROPS

The presence of numerous skin wild species and ancestres of domesticated plants are an indicator of the diversity that may be expected from the Andes. Those crops which have a high priority in the Andean Region due to their greater proliferation, appropriate nutritional value and efficient reproduction system, are listed in Table 3.

T	A	3L	ε	3

MAIN ANDEAN NATIVE CROPS IN PERU

.

	Crop Area	Altitude
	in ha	mas 1
		-
GRAINS		
Quinoa l	15,000	0-3,900
Kaniwa	5,000	3,900-4,100
Amaranthus	500	0-3,000
Lupinus	3,000	1,500-3,800
TUBERS		
Oca	. 18,000	3,000-4,000
Olluco	16,000 .	2,000-3,900
Mashua	6, 000	2,800-4,000
ROOTS		
Yacón	300	0-3,000
Arracacha	400	0-3,000

I It is estimated that there are 40,000 ha of quinoa in Bolivia and 2,000 ha in Ecuador.

_~?

-

-

~

ه سی د ی بور....

Quinoa (<u>Chenopodium quinoa</u>) is generally found in the Andean highlands from 3,500 to 4,000 m of altitude. However, it has been established also that it grows from sea level to near the snow caps.

TABLE 4

QUINOA MAIN GROUPS AND THEIR ECOLOGICAL ADAPTABILITY

-

	GROUP	ALTITUDE	COUNTRY/REGION
-	Sea level Quinoa .	0- 500	Chile/Central Zone
-	Valley's Quinoa	1,000-3,000	Perú, Bolivia, Ecuador, Colom bia
-	Highland Plains Quinoa	3,500-33,900	Per ú, Bolivia
-	"Salares" Quinoa	3,700-3,800	Bolivia
-	High "Yungas" Quinoa	2,500-3,000	Bolivia/Perú

Gandarillas (1968) describes eighteen races, classified as per their growth habitat, type of inflorescence and shape and edge of leaves.

Quinoa's configuration ranges from branches to one single stem plants and shows amaranth or glomerulous leaf types and smooth to saw-edge leaves. The height of this plant ranges from 1 m to 3 m. The smallest ecotypes are found in the Highland plains and the higher ecotypes in the inter-Andean valleys located at 3,000 masl.

Given this broad diversity, it is possible to select some varieties which can endure low humidity. Quinoas may be grown in salty pans ("salares") with only 280 mm of rain provided that specific agricultural techniques, which consist of small holes dugged in the ground, are used. Quinoas grown in the valleys may withstand rainfalls over 1,000 mm. Tolerance to fungus diseases, such as mildiu (<u>Peronospora</u> sp), is associated with rainfall and humidity.

The colour of the grain varies from dark (usually associated with a higher protein content, up to 21%) to white. Grains can be amilaceous as well as crystalline ("chullpi") ranging from 12 to 16% protein. Additionally, there is a group of gray skinned quinoas called "ccoito" which produce soft flour.

The adaptability of quinoas to different soils is related to their pH. The "royal quinoas" grown in the "salares" are highly halophyte and can grow in alkaline soils (pH 7.8). The sea level quinoas show good adaptability to neutral pH's.

Kañiwa is possibly the most resistant crop to low temperatures existing in the world. This plant has a special anatomical structure which protects the inflorescence and can withstand temperatures as low as -3°C during flowering.

Its grains show different hues ranging from light brown to black. The grain cover (perisperm) ranges from light gray, yellow, orange or dark red to almost black.

The Andean Amaranthus is identified by way of two species : <u>A. caudatus</u> in the Central Andean mountains in Perú and Bolivia and <u>A. edulis</u> in Southern Bolivia and Northern Argentina, known under names such as "coyo", "achis", "achita", "coimi", and "millmi", which often lead to confusion. A similar specie is <u>A. hibridus</u> known as "hat'ago" eaten in its tender stage as a vegetable. The height of this plant ranges between 1 m and 3 m.

Tarwi or chocho (Lupinus mutabilis) also shows a broad diversity. Three different types may be clearly identified, as indicated in Table 5.

TABLE 5

			· · ·
COMMON NAME	AREA	ALTITUDE	GROWTH
Tarwi or tauri	Bolivia, Southern Perú	3,600-3,800	Slightly ramified Semi-early
Chocho	Central and Northern Peru, Ecuador	3,000-	Ramified/late type
Tarwi	Around Lake Titicaca	3,800-3,850	Slightly ramified/ very early (150 days)

DIVERSITY OF LUPINUS MUTABILIS

Several authors mention the ample variability of Lupinus wild species. McBride (1953) indicates the existence of 82 species, some of which are grown and used as food. Carrillo (1947) has described most of these species, one of which is <u>Lupinus praestabilis</u> grown in altitudes up to 4,200 masl. Tapia (1980) has suggested the kind of germplasm that could be used in a breeding programme for the enhancement of man-grown species, with emphasis on resistance to frost, diseases and pests.

TUBERS

Andean tubers such as Oca (Oxalis tuberosa), Olluco (Ullucus tuberosus) and Mashua or Isaño (Tropaeolum tuberosum) do not vary only in their shape, size and color of tuber, but also in their precocity, tolerance to cold, and oxalic acid content, as in the case of oca, and mucilagous content as in the case of olluco (Table 6).

This broad genetic diversity, a common indicator of the Andean species, should be used advantageously in a cooperative programme among countries having high mountain conditions, in order to obtain improved materials adapted to these environments.

Diseases and pests are to be considered within the framework of a plan for the selection of tolerant ecotypes and should be a logical alternative to the use of chemical products.

TABLE 6

CHARACTERISTICS OF OCA TUBERS 1

COMMON NAME	ORIGIN	COLOUR	LENGHT OF STYLE
Yurac Oca	Bolivia, Potosí	White	Long
Pallihuaya	Bolivia, Omasuyo	Yellow	Short
Puca Oca	Bolivia, Cochabamba	Red	Short
Keni Pecke Oca	Bolivia, Pillapi	White	Medium
Ibia	Colombia, Zipaquirá	Light Yellow	Medium
Lari Oca	Bolivia, Tunari	Purple	Medium

1 Source : Cárdenas (1958)

Generally speaking, resistance to low temperatures is associated with height of plants. Normally, species growing very close to the ground surface would withstand better lower temperatures.

R00T5

Yacon is grown from Venezuela to Argentina. It is a perennial root, even though aerial stems are annual. These roots are eaten raw and have a sweetish flavour (Leon, 1964).

Arracacha is a species considered to be the oldest in America (Bukasov, 1930). As it is a short photoperiodic plant, its introduction to places outside the tropics has been impossible. (Montaldo, 1972).

There are white and purple varieties adaptable up to 2,700 m. Plants produce 4-10 cone shaped roots.

ANIMALS

SOUTH AMERICAN CAMELIDS

Out of the four camelids found in the Andean region, two of them, the llama and elpaca are domesticated.

These two animals fulfill different though complementary functions. The llames are taller and differ in height: adult llamas found in the Bolivian "salares" are 2.20 m tall and those found along the drier high desertic lands ("punas") are 1.60 m tall. Llamas are used as means of transportation of agricultural products (potato, barley and fruits) along the highlands. Their fiber though somewhat thicker, is highly appreciated for its strength and durability. Meat of younger animals is widely accepted and a variety of products is produced with the leather, such as clothing and straps for agricultural and construction products. There are three distinctive types: two with a lot of fiber which differ according to their height, and a third group of animals almost hairless.

Alpacas are finer and have a higher quality of fiber. An adult alpaca produces 2,103 kg per year. Alpaca meat is highly appreciated and it has been widely used in the sausage industry.

TABLE 7

COUNTRY	ALPACAS	LLAMAS	TOTAL
Perú	3'290,000	900,000	4 '190,0 00
Bolivia	300,000	2'000,000	2'309,000
Chile	50, 000	70,000	120,000
Argentina	Few	Few	-
TOTAL	3'640,000	2'970,000	6'610 ,0 00

CAMELIDS POPULATION IN SOUTH AMERICA, FAD (1971)

GUINEA PIG "CUY"

In Perú, at present, there are about 20'000,000 "cuys" born per year (Arroyo, 1986), and the annual consumption is estimated in 1.2 kg per capita. They are bred by families, and there are four different species.

A research study carried out by Calero del Mar (1978) on the variability of these species, shows that their hair varies from white to black, and their body shape is adequate for beef production, except those with a more angular type. The amount of hair is also highly variable.

TABLE 8

COMPARISON OF CONVERSION EFFICIENCY BETWEEN GUINEA PIG AND OTHER SPECIES (ALIAGA, 1979)

WEIGHT/KG	CONSUMPTION PER WEIGHT	INCREASE/WEIGHT
	x	%
an a	****	·
0.8	31.25	0.9
40.0	12.50	0.3
500.0	10.0	0.2
	WEIGHT/KG 0.8 40.0 500.0	WEIGHT/KG CONSUMPTION PER WEIGHT % 0.8 31.25 40.0 12.50 500.0 10.0

TABLE 9

ANNUAL PRODUCTION OF MAIN DOMESTICATED ANDEAN SPECIES

	Meat/kg/unit	Fiber kg	Milk/l/day
Llamas	40 - 60	3 - 4	1 - 3
Alpacas	30 - 50	2 - 5	1 - 2
Guinea Pig	0.3 - 0.8	-	-

· •

III. COLLECTION EXPEDITIONS : THE ANDEAN NATIVE CROPS AND DOMESTIC ANIMAL RESOURCES GERMPLASM BANKS.

Numerous isolated efforts have been deployed in order to collect this material. Probably one of the most relevant expeditions was the one compiled by Dr. Jorge Leon, and Eng. Julio Rea on the early 60's, when the Programa de Cultivos Andinos of IICA was first started. These collections included mainly tubers and roots.

The material was distributed to the universities of Ecuador, Perú, and Bolivia, and a variable percentage of this material is still available. A large portion of the material on oca is kept in Cusco and Huancayo, and the germplasm on arracacha is kept at the Cajamarca University in Perú.

The biggest collection was sent to Dr. Martín Cárdenas of Cochabamba University in Bolivia. Upon the establishment of accessions, important evaluations were made, and publications were produced (Cárdenas, 1958).

During the late 60's and early 70's several expeditions were made by the Peruvian regional universities giving special importance to Andean grains.

In Puno, most efforts were devoted to Quinoa and Kañiwa, while in Cusco and Huancayo, Lupinus was the main priority.

As of 1975, a coordinating action for the organization and compilation of the collections existing in Bolivia and Perú, was initiated under the Program Andes Altos of IICA, based in Bolivia. Support was also provided to similar actions undertaken in Colombia and Ecuador.

This program was further supported, founded and guided by FAD's International Committee on Phytogenetic Resources (CIRF).

Apart from the support received for several expeditions between 1976-1980, the following facilities were constructed and/or endowed :

Bolivia

Belén Experimental Station (E.S). Improvement and endowement of installed facilities.

Patacamaya E.S. Purchase of equipment.

Perú

Camacani E.S. (Puno). Building and equipment. Kayra E.S. (Cusco). Building and equipment. E.S. (Ayacucho). Building of facilities.

The setting of this infrastructure facilitated the identification of the material, as well as the selection of improved lines.

The Lupinus project supported by GTZ, was carried out in Perú during eight years, and contributed to the collection of these species. The above mentioned germplasm was distributed between Cusco and Huancayo and the construction of an appropriate building was funded.

Up to the present, CIRF and IDRC have continued supporting the maintenance and evaluation of specific collections in Ecuador, Perú, and Bolivia.

In 1982, the Amaranth collection (227 accessions) was given to Cusco University leading to the establishment of a program for the promotion of this crop.

In 1985, the Instituto Nacional de Investigación y Promoción Agropecuaria del Perú (INIPA), created the Programa Nacional de Cultivos Andinos (National Andean Crops Program) which is now concentrating all efforts on the research of these species.

. •

TABLE 10

INSTITUTIONS RESPONSIBLE FOR GERMPLASM

•

CROP	INSTITUTION
Quinoa-Kañiwa	- CIPA XXI-PUNO, c/o Ing. Valeriano Huanco Apartado 388 Puno, Perú
	- UNA, c/o Ing. Roberto Valdivia (Universidad Nacional del Altiplano) Apartado 271 Puno, Perú
Lupinus	 Universidad del Cusco, c/o Ing. Oscar Blanco Apartado 1006 Cusco, Perú
	 CIPA XVI-JUNIN, c/o Ing. Oscar Garay Calle Real 509 El Tambo, Huancayo, Perú
Andean Tubers and Kiwicha	- Universidad del Cusco, c/o Ings. Hernán Cortéz/Luis Sumar Apartado 1006 Cusco, Perú
	- Universidad de Ayacucho c/o Ing. Julio Valladolid Apartado 243 Ayacucho, Perú
Andean Crops/General	 Programa Nacional de Cultivos Andinos c/o Dr. Mario Tapia Apartado 110097 Telex 25194 NC Lima 11, Perú

- 16 -

.

TABLE 11

INSTITUTIONS RESPONSIBLE FOR THE ANDEAN ANIMAL GERMPLASM

ANIMAL	INSTITUTION
Camelids	- IVITA, Universidad Nacional Mayor de San Marcos Apartado 4270 Lima, Perú
	- INIPA, c/o Ing. Julio Sumar Apartado 110097 Lima 11, Perú
Guinea Pigs	- INIPA, c/o Ing. Marco Saldivar Apartado 110097 Lima 11, Perú

.

,

۰

•

ÎN Î	VIBER OF ACCESSIONS	(COLLECTIONS) PER	CROP IN EACH GERMPI	LASM BANK (1985)	IN PERU
	PUNO	cusco	HUANCAYO	АҮАСИСНО	CAJAHARCA
Quinoa	1500 (E)	198	48	425	425
Kañiwa	330	,	14	47	ı
Kiwicha	ı	570	32	109	17
Tarwi	228 (E)	'1200 (E)	1500(E)	325	126
Bitter Potato .	68	130	. 42	257	ı
Oca	120	610	168	122	30
011uco	40	13	118	61	18
Isaño, Mashua	65	. 14	47	107	t

,

TABLE 12

(E) Include material from other Andean countries.

IV. PRESENT AND POTENTIAL USE OF ANDEAN PHYTOGENETIC AND ZOOGENETIC RESOURCES

Andean crops, other than potatoes and corn, have not been used appropriately. It is a result of the food crisis emerged in this decade which has reached critical levels, (Perú imports more than 90% of the wheat and 30% of the milk consumed), that the value of these species has been recognized. They should be suitably analized, improved and promoted as they may prove to be more advantageous than many imported food products.

The surface area of the main crops mentioned here, occupies at present from 60,000 to 70,000 ha; yields and prices vary according to the year and the prevailing climatic conditions (Table 11).

Andean grains such as Quinoa and Kiwicha have awakened great interest. However, their nutritional role has not been clearly defined as yet despite the fact that peasants have traditionally used them for soups and sweet and salty dishes replacing rice.

The grains of Quinoa and Kiwicha have been processed industrially to produce flour which can be mixed with milk, or used to prepare "tortillas" (flat bread). Pop Quinoa, and Kiwicha are also common.

Although Kañiwa will eventually cover a large area (over 50,000 ha), it requires to be roasted before consumption. Its nutritional value, of a high protein content (15%), has been widely recognized.

Tarwi is also rich in proteins (32-42%). It is a legume considered as the Andean soy bean. If processed to release its alkaloid content, it may be used as a food product similar to beans.

The use of this legume in rotations may substantially increase the production of other crops, as it is estimated to fix between 60 to 80 kg of N/ha.

Roots and tubers are an important source of energy. Their future perspective will improve when industrial processing of the flour gets under way.

One hectare of Oca may produce up to 8 T of dry matter/ha, thus competing favourably with other cereals.

Common agronomic practices for these crops are shown in Tables 12 and 13.

Herds of South American camelids such as llama and alpaca reached more than 20'000,000, approximately three times their present population. They were an important source of proteins for the population of the XVI century when the Spaniards arrived.

The selection of animals used to be done yearly in big rodeos "chaco" which permitted choosing the surplus and adult animals. Meat was dried and salted under the sunlight to preserve it for long periods of time.

The above technique is still practiced and allows meat to be transported to the cropical region. Given the fact that these species graze easily in the Andean pastures, their population could increase notoriously if competition by cattle and sheep is reduced (Primov, 1983).

Alpaca hair is highly appreciated in the world markets and countries such as Bolivia and Perú still retain more than 95% of the world's total production.

Guinea pig raising is the most economical source of proteins, because all residues from the kitchen may be used for their feeding. Some selected guinea pigs have reached 1,5 kg in weight.

Guinea pigs may be cooked in different ways such as broiled or fried. Traditionally, it is served on holidays.

As indicated in the opening statement of this article, it is important to know the geography and the resources available in a given region because it allows for the development of agricultural practices and crops which may suit better the prevalent mountain conditions. These resources may be shared, as in the case of potatoes and corn, with other mountaneous countries of the world.

	DIST	RIBUTION OF AN	DEAN CROPS, A	LTITUDE, AND	VIELDS IN TH	E PERUVIAN HI	GHLANDS	
		CAJAMARCA LA LIBERTAD	ANCASH	VALLE DE MANTARO	AYACUCHO APURIMAC	cusco	PUNO	TOTAL AREA ' HA
QUINOA	0 K K D	** 2600-3500 1500	* 2700-3800 1000	* 2500-3700 1400	** 2600-3800 800		*** *** 3800-3900 600-1500	16,000
KANIWA	٥	;	;	*	*	**	***	4,000
KIWICHA	Ο	*	- *	*	· *	*		800
TARWI .	047	** 2000-3300 1000	** 2000-3400 800	* 2500-3500 800-2000	* 2200-3500 900	** 2400-3500 800-2000	** 3800 1000-2500	5,000
BITTER POTATO	047	9 8	;	* 3800 8000	* 3800 - 4000 7000	** 3800-4000 7000	** 3900-4000 7000	38,000
0CA (2)	04	** 3000-380U	** 3000-3800	** 3000-3900	** 3000-	* *	*	16,000
OLLUCO	Δ, A D	** 3000-3800	** 3000-3900	** 3000 -3 800	* *	*	¥	10,000
ISAÑO, MASHUA	047	* 3000-3800	* 3000-3500 7	* 3000-3900	10000 10000	*	*	6,000
D = Distributior		A = Altitude i	n masl	Y = Yield P	, cg/ha (1)			
<pre>(1) Yields vary (2) Andean tuber</pre>	due 's are	to technologi e normally int	cal levels, y ercropped or	/ears and var associated w	iety used. ith maize, th	us yields are	e not include	ed in all

TABLE 13

cases.

	ACRONUMIC	PRACTICES ON ANDEAN GRA	SNI	
ACTIVITY	KIWICHA	VONIND	KANIWA	TARWI
Planting Uale Method Seed Treatment Seeding rate kg/ha Planting distance cm	August-October Rows N.N 3-6 60-80	August-November Row-Transplanting N.N 4-15 40-80	September-October Row-Broadcast N.N 3-6 40-50	August-October Row-/In holes Fungicide 60-80 60-80
Fertilization	80-40-0	80-40-0	60-40-0	0-40-0
Hilling Up	At least 1	1-2	N.N.	-1
Control Diseases	Esclerotinea Alternaria	Peronospora Ascochyta Phoma Pseudomonas		Colletrotrichum Uromyces, Chrysocelis
Pests	S. Z	Ticona Copitarsia Scrobipalpula Pachyzancla Epicauta Myzus Frankliniella Astillus	۲ ۲	Agromyza Apion
Birds	N.N	During last 3 months	N.N.	N . N
Harvest Method	Plant cutting	Plant cutting or uproot	Uproot or cut	Plant cutting
Threshing	Manual or . mochanical	Manual or mechanical	Manual	Manual
Grain Classification	Wind cleaning or mechanical	Wind cleaning or mechanical	Wind cleaning	By hand
<pre>N_Significant "Not Needed</pre>				

TABLE 14

MASHUA	1000-1200	60-80	100-60-0	2-3	"Fasciación"**		220-245	N.S
OLLUCO	800-200	60-80	60-40-0	2-3	"Fasciación"**	Premnotrypex	210-245	N.S
OCA	1200-1400	60-80	120-20-0	2-3	N.S	Copitarsia Bothynus Macrosiphum	220-270	Rotting Rhizopus
ACTIVITY	Seeding rate kg/ha	Planting distance cm	Fertilization *	Hilling-up (number)	Diseases	Pests	Harvest (Vegetative cycle)	Storage (problems)

r

Fertilization is conditioned to the rotational system. In new fields rates are very low, less after fertilized potato and higher after a cereal. The use of manure is common but the rates used are very low. ¥

•

•

** Physiological disorder, undefined origin.

N.S. = Non significant

TABLE 15

AGRONOMIC PRACTICES - ANDEAN TUBERS

.

BIBLIOGRAPHY

.

- ALIAGA, L., 1979. Producción de cuyes. Huancayo, Universidad Nacional del Centro.
- ARROYO, O., 1986. Investigaciones sobre el cuy en el Perú. Proyecto PISA-INIPA-CIID-ACDI. Informes Técnicos No. 8. Lima, Perú.
- BUKASOV, S.M., 1930. The cultivated plants of Mexico, Guatemala and Colombia. Bull. Appl. Bot. Genetic Plant Breeding. Supp. 47

CALERO DEL MAR, B., 1978. El cuy. Ed. Agronómica. Cusco, Perú. 281 p.

- CARDENAS, M., 1958. II Informe sobre trabajos hechos en Bolivia sobre Oca, Olluco y Mashua. En : Estudios sobre tubérculos alimenticios de los Andes. Comunicaciones de Turrialba No. 63. Costa Rica.
- CARDOZO, A., 1954. Auquénidos. La Paz, Bolivia.
- CARRILLO, E., 1956. Revisión del género Lupinus en el Perú. Tesis Universidad Nacional de San Agustín. Arequipa, Perú.
- FAO, 1971. Estadísticas de ganadería. Yearbook, Rome.
- GANDARILLAS, H., 1968. Razas de Quinua. Boletín experimental No. 34. Ministerio de Agricultura. La Paz, Bolivia.
- HORKHEIMER, H., 1973. Alimentación y obtención de alimentos en el Perú prehispánico. UNMSM. Lima, Perú.
- LEON, J., 1964. Plantas alimenticias andinas. Boletín Técnico No. 6. IICA. Lima, Perú.
- MC. BRIDE, J.F., 1953. Flora of Perú. Field Museum of Natural History. Botanical Series 13. Chicago, 510 p.
- MONTALDO, A., 1972. Cultivo de raíces y tubérculos tropicales. IICA. Lima, Perú.
- PRIMOV, G., 1983. Alpaca meat production and exchange in Southern Prú. Technical report series No. 31. SR/CRSP. Lima, Perú.
- TAPIA, M.C. VARGAS, 1982. Wild lupine of the Andes of Southern Perú. Agricultural and nutritional aspects of lupines. Proceedings of the First International Lupine Workshop. Lima-Cusco, Perú.
- TAPIA, M.E. et al., 1980. La Quinua y Kañiwa. Cultivos andinos. Pub. IICA. Bogotá, Colombia.

- VAVILOV, N.I., 1935. The origin, variation inmunity and breeding of cultivated plants. (English translation). Chronica Botanica Co. Waltham, Mass., USA.
- WING, E.S., 1975. La domesticación de animales en los Andes. Rev. Allpanchis No. 8. Cusco, Perú.
- YACOLEFF, E.; HERRERA, F.L., 1934. El mundo vegetal de los antiguos peruanos, Lima. Rev. Museo Nacional. Tomo III No. 3. 241-323 p.













- FIG.2 THE ANDEAN GRAINS
 - Quinoa Kañiwa 1.
 - 2.
 - 3. Kiwicha or Amarantus Tarwi or Lupinus
 - 4.





FIG. 3 THE ANDEAN TUBERS

-

1. Olluco 2. Oca 3. Isaño or mashua







FIG. 5 THE SOUTH AMERICAN CAMELIDS (Cardozo, 1954)



FIG. 6 DISTRIBUTION OF THE CAMELIDS