

# JORDAN STILLBIRTH AND NEONATAL MORTALITY SURVEILLANCE (JSANDS) SYSTEM

©2024,



This work is licensed under the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/legalcode>), which permits unrestricted use, distribution, and reproduction, provided the original work is properly credited. Cette œuvre est mise à disposition selon les termes de la licence Creative Commons Attribution (<https://creativecommons.org/licenses/by/4.0/legalcode>), qui permet l'utilisation, la distribution et la reproduction sans restriction, pourvu que le mérite de la création originale soit adéquatement reconnu.

*IDRC GRANT / SUBVENTION DU CRDI : - IMPLEMENTING A PERINATAL AND NEONATAL MORTALITY SURVEILLANCE AND AUDITING SYSTEM IN JORDAN*



# Disparities in Obstetric, Neonatal, and Birth Outcomes Among Syrian Women Refugees and Jordanian Women

Tariq N. Al-Shatanawi<sup>1</sup>, Yousef Khader<sup>2</sup>, Nadin Abdel Razeq<sup>3\*</sup>, Ahmed M. Khader<sup>4</sup>, Mahmoud Alfaqih<sup>5</sup>, Osama Alkouri<sup>6</sup> and Mohammad Alyahya<sup>7</sup>

<sup>1</sup>Department of Public Health and Community Medicine, Al-Balqa Applied University, Al-Salt, Jordan, <sup>2</sup>Department of Public Health, Jordan University of Science and Technology, Irbid, Jordan, <sup>3</sup>Department of Maternal and Child Health Nursing, School of Nursing, The University of Jordan, Amman, Jordan, <sup>4</sup>Faculty of Medicine, Jordan University of Science and Technology, Irbid, Jordan, <sup>5</sup>Department of Physiology and Biochemistry, Jordan University of Science and Technology, Irbid, Jordan, <sup>6</sup>Faculty of Nursing, Yarmouk University, Irbid, Jordan, <sup>7</sup>Department of Health Management and Policy, Jordan University of Science and Technology, Irbid, Jordan

**Objectives:** To compare obstetric and neonatal characteristics and birth outcomes between Syrian refugees and native women in Jordan.

**Methods:** We used the Jordan Stillbirths and Neonatal Deaths Surveillance System to extract sociodemographic and obstetric characteristics of the mothers and birth characteristics of newborns. Multivariate analysis was used to compare the characteristics of 26,139 Jordanian women (27,468 births) and 3,453 Syrian women refugees (3,638 births) who gave birth in five referral hospitals (May 2019 and December 2020).

**Results:** The proportions of low birthweight (14.1% vs. 11.8%,  $p < 0.001$ ) and small for gestational age (12.0% vs. 10.0%,  $p < 0.001$ ) newborns were significantly higher for those born to Syrian women compared to those born to Jordanian women. The stillbirth rate (15.1 vs. 9.9 per 1,000 births,  $p = 0.003$ ), the neonatal death rate (21.2 vs. 13.2 per 1,000 live births,  $p < 0.001$ ), and perinatal death rate (21.2 vs. 13.2 per 1,000 births,  $p < 0.001$ ) were significantly higher for the Syrian births. After adjusting for sociodemographic and obstetric characteristics of women, only perinatal death was statistically significantly higher among Syrian babies compared to Jordanian babies (OR = 1.3, 95% CI: 1.1–1.7,  $p = 0.035$ ).

**Conclusion:** Syrian refugee mothers had a significantly higher risk of adverse obstetric and neonatal outcomes including higher rate of perinatal death compared to Jordanian women.

**Keywords:** Jordan, mortality, Syrian refugee, pregnancy outcomes, neonates

## OPEN ACCESS

### Edited by:

Andrea Madarasova Geckova,  
University of Pavol Jozef Šafárik,  
Slovakia

### Reviewed by:

Ankit Anand,  
Institute for Social and Economic  
Change, India  
One reviewer who chose to remain  
anonymous

### \*Correspondence:

Nadin Abdel Razeq  
n.abdelrazeq@ju.edu.jo

**Received:** 01 December 2022

**Accepted:** 19 October 2023

**Published:** 03 November 2023

### Citation:

Al-Shatanawi TN, Khader Y,  
Abdel Razeq N, Khader AM, Alfaqih M,  
Alkouri O and Alyahya M (2023)  
Disparities in Obstetric, Neonatal, and  
Birth Outcomes Among Syrian Women  
Refugees and Jordanian Women.  
*Int J Public Health* 68:1605645.  
doi: 10.3389/ijph.2023.1605645

## INTRODUCTION

Forced migration occurs globally for various reasons, including wars and political conflicts, economic circumstances, or environmental crises [1]. Women refugees of all nationalities are a particularly vulnerable group of displaced people. Having a higher reported prevalence of adverse maternal-neonatal outcomes than native women [2, 3] makes them a unique population in host communities in terms of needed humanitarian and healthcare services. Stillbirths, delayed arrival at

the hospital for birth [4, 5], inappropriate weight for gestational age [2], low birthweight infants [6], and higher rates of neonatal intensive treatments for neonatal comorbidities [2] are a few adverse pregnancy outcomes that are commonly reported among women refugees. However, contemporary research and knowledge on women refugees' health during pregnancy and pregnancy outcomes is limited in the international literature. Therefore, further understanding of how their health is compared to non-refugee counterparts is needed [3].

Since March 2011, Syria has been in a state of political instability that has resulted in about 13.5 million Syrians being forcibly displaced to escape the war, out of which 6.8 million sought refuge in their country [7]. The neighbouring country of Jordan has been the host of more than one million Syrian refugees [8], of which only 673,957 have registered with the United Nations High Commissioner for Refugees (UNHCR) as of February 2022 [9]. Women and girls constitute 49.6% of all registered refugees in Jordan, and 30% of them are above the age of 12 years [9]. Further, reports indicate that the majority (>85%) of registered Syrian refugees live outside camps, in urban, peri-urban, or rural regions of Jordan [9]. According to recent assessments of the needs of refugees for financial and non-financial services [10, 11], more than 86% of Syrian refugees live under the Jordanian poverty line [11] despite the substantial response by government and humanitarian partners.

In Jordan, healthcare services, including pregnancy and delivery services, are available to all registered refugees from all nationalities at the non-insured Jordanian rate at public health centres and governmental hospitals [12]. However, because most Syrian refugees are not insured and must pay health-related fees for hospitals and clinics, Syrian refugees identified cost as the main barrier to healthcare access [13]. The UNHCR supports healthcare services for all refugees. However, research has shown that Syrian women refugees in Jordan continue to have significant unmet needs and barriers to access, utilisation, and implementation of sexual and reproductive health (SRH) in comparison to their host community counterparts [14, 15].

There is limited research assessing the reproductive health status and service delivery among Syrian refugees in Jordan [14]. Currently, there is little understanding of the risks of adverse pregnancy outcomes among Syrian women refugees in Jordan and the extent to which disparities exist regarding pregnancy outcomes. The adverse maternal and neonatal pregnancy outcomes and disparities need to be addressed first, and their contributing factors should be systematically investigated to improve the maternal healthcare utilisation and health services provided to Syrian women refugees.

The Jordan Stillbirths and Neonatal Deaths Surveillance System (JSANDS) is an electronically secure surveillance system to register and disseminate reliable, comprehensive, individual-level data on stillbirths and neonatal deaths, the causes and the risk factors most commonly associated with perinatal mortality in Jordan [16–18]. The system record events for women prospectively after giving their births. The surveillance system automatically and accurately transfers the data on births, stillbirths, and neonatal deaths from five maternity hospitals to the Ministry of Health [16]. These hospitals are

among the most significant and primary referral hospitals in Jordan that cover most births and deaths in all regions in the north, east, and south of Jordan. Khader, Alyahya [16] examined the JSANDS attributes, including its data quality and sensitivity. The system users rated the usefulness of JSANDS as excellent (percentage score = 85.6%). The overall acceptability (percentage score = 82.3%), flexibility (percentage score = 80.2%), stability (percentage score = 80.0%), and representativeness (percentage score = 86.6%) were also rated excellent. All variables in JSANDS had complete data with no missing values for clinical variables because these data are entered by nurses who were trained to enter data completely and because the system has automated quality checks. The overall simplicity was scored well (percentage score = 75.4%) [16].

The purpose of this study is to explore the adverse birth outcomes among Syrian women refugees and Jordanian women using data from the JSANDS. Specifically, in this study we identified disparities of the obstetric and neonatal characteristics and birth outcomes between Syrian and Jordanian women, and explored the factors determining adverse neonatal birth outcomes among Syrian women in Jordan.

## METHODS

### Study Design

All births, stillbirths, and neonatal deaths in five hospitals from May 2019 to December 2020 were completely registered in JSANDS. The five hospitals, including one university teaching hospital, three public hospitals, and one private hospital, were selected for pilot testing of the JSANDS. The criteria for the selection of hospitals and JSANDS usability and functionality were described in previous studies [17, 19, 20]. The sociodemographic and obstetric characteristics of Jordanian and Syrian women and their birth outcomes were retrieved from JSANDS and compared.

### Variables

The extracted data consisted of sociodemographic and obstetric characteristics of the mothers, including age, education, employment status, income, gestational age, mode of delivery, and multiplicity. Other data included information on neonatal characteristics and birth outcomes, such as birthweight and status (born alive, stillbirth, and neonatal death).

### Definitions

Birthweight was categorised into low (1,500 g–<2,500 g), very low (<1,500 g–1,000 g), and extremely low birthweight (<1,000 g) [21]. The gestational age was categorised into extremely preterm (<28 weeks), very preterm (28–<32 weeks), and moderate or late preterm (32–<37 completed weeks of gestation) [22]. The definition of stillbirths and neonatal deaths used in this study was based on the international standards set by the World Health Organization [23]. Neonatal mortality was defined as any death that happened at or after 24 gestational weeks within the first 28 days of life. The neonatal mortality rate was calculated as the number of neonatal deaths per 1,000 live births (LB). Stillbirth was

**TABLE 1** | The sociodemographic and obstetric characteristics of Jordanian and Syrian women (Jordan, May 2019 and December 2020).

	Nationality				p-value
	Jordanian		Syrian refugees		
	n	%	n	%	
Mother's age (years)					<0.001
<20	982	3.8	423	12.3	
20–35	20,590	78.8	2,590	75.0	
>35	4,567	17.5	440	12.7	
Mother's occupation					<0.001
Housewife	23,252	89.0	3,397	98.4	
Employed	2,887	11.0	56	1.6	
Mother's education					<0.001
Diploma or less	16,713	68.6	2,290	94.7	
Bachelor or higher	7,660	31.4	129	5.3	
Income (JD)					<0.001
≤500	19,901	80.5	3,087	96.7	
>500	4,811	19.5	105	3.3	
Type of hospital at birth					<0.001
Public hospital	17,376	66.5	2,376	68.8	
Private hospital	4,872	18.6	1,036	30.0	
Teaching hospital	3,891	14.9	41	1.2	
Multiplicity					0.022
Singleton	24,916	95.3	3,299	95.5	
Twin	1,140	4.4	134	3.9	
Triplet or more	83	0.3	20	0.6	
Parity					<0.001
Nulliparity	842	3.2	92	2.7	
Low multiparity (parity 1–3)	17,414	66.6	1,992	57.7	
Grand multipara (parity ≥4)	7,883	30.2	1,369	39.6	
History of neonatal death	447	1.7	85	2.5	0.002
History of stillbirth	4,935	18.9	652	18.9	0.997
Gestational age					0.137
Full-term	23,761	90.9	3,112	90.1	
Preterm	2,378	9.1	341	9.9	
Mode of delivery					<0.001
Vaginal	13,017	49.8	1,772	51.3	
Planned CS	6,982	26.7	1,060	30.7	
Emergency CS	5,957	22.8	604	17.5	
Instrumental	183	0.7	17	0.5	

defined as any foetal death that occurred at or after 24 gestational weeks. Stillbirths were categorised as antepartum (deaths occurring prior to labour) and intrapartum (deaths that occur after the onset of labour but prior to birth). The stillbirth rate was calculated as the number of stillbirths per 1,000 total births. The foetal weight percentile was determined based on birthweight and foetal age. Foetal growth was categorised as small for gestational age (SGA) if <10th percentile, appropriate for gestation if between 10th and 90th percentile, or large for gestation if >90th percentile.

## Ethical Considerations

This study was performed in line with the principles of the Declaration of Helsinki. The study was ethically approved by Jordan University of Science and Technology Institutional

Review Board (IRB) (Ethical approval number 20170033) to conduct the study at King Abdulla University Hospital, Jordan Ministry of Health Review Committee to conduct the study in the three public hospitals, and the Irbid Speciality Hospital (private hospital) Ethical Committee. Permission from the Ministry of Health was received to access the data and use it for research purposes. To ensure the confidentiality of the data, the data were exported without identifying information, such as the name or phone number.

## Data Analysis

Data were described and analysed using IBM SPSS version 24 (IBM Corp. Released 2016. IBM SPSS Statistics for Windows, Version 24.0. Armonk, NY: IBM Corp.). Percentages were used for categorical data. The sociodemographic and obstetric characteristics of Jordanian and Syrian women and their birth outcomes were compared and analysed using the Chi-square test. Multivariate analysis using binary logistic regression was performed to compare the birth outcomes of Jordanian and Syrian women after adjusting for women's characteristics. A *p*-value of less than 0.05 was considered statistically significant.

## RESULTS

### Women's Characteristics

This study included a total of 26,139 Jordanian women (27,468 births) and 3,453 Syrian women refugees (3,638 births). Of the total Jordanian and Syrian births, 271 and 55 were stillbirths, and 360 and 76 died before 28 days of life, respectively. The sociodemographic and obstetric characteristics of Jordanian and Syrian women are shown and compared in **Table 1**.



Syrian women were statistically significantly younger than Jordanian women (12% and 4% under age 20 years, respectively, *p* < 0.001). The mean (SD) age was 29.3 (6.0) year for Jordanian mothers and 27.3 (6.7) year for Syrian women. Only 11.0% and 1.6% of Jordanian and Syrian women were employed, respectively (*p* < 0.001). Jordanian women had a higher level of education than Syrian women (31.4% and 5.3% of Jordan and Syrian women had bachelor's or higher degrees, respectively). Moreover, monthly income was statistically significantly higher for Jordanian women than for Syrian women. Jordanian and Syrian women differed statistically significantly in terms of multiplicity rate, parity, and mode of delivery. The mean (SD) parity was 2.8 (1.7) for Jordanian mothers and 3.3 (2.0) year for Syrian women.

### Birth Outcomes

**Table 2** shows the birth outcomes for Jordanian and Syrian newborns. In comparison to the babies born to Jordanian women, the proportion of babies born with low weight was statistically significantly higher for Syrian women (11.8% vs. 14.1%, *p* < 0.001). About 10.0% and 12.0% of babies born to Jordanian and Syrian women were SGA (*p* < 0.001), respectively.

**TABLE 2** | Birth outcomes for Jordanian and Syrian newborns (Jordan, May 2019 and December 2020).

	Nationality				<i>p</i> -value
	Jordanian		Syrian refugees		
	<i>n</i>	%	<i>n</i>	%	
Preterm birth	2,874	10.5	415	11.4	0.082
Low birthweight	3,236	11.8	512	14.1	<0.001
Small for GA	2,742	10.0	437	12.0	<0.001
Stillbirth rate (per 1,000 total births)	271	9.9	55	15.1	0.003
Neonatal death rate (per 1,000 live births)	360	13.2	76	21.2	<0.001
Perinatal death rate (per 1,000 total births)	549	20.0	116	31.9	<0.001




The distribution of preterm delivery and birthweight for Syrian and Jordanian babies is shown in **Figures 1, 2**, respectively. The stillbirth rate (15.1 vs. 9.9 per 1,000 total births,  $p = 0.003$ ), the neonatal death rate (21.2 vs. 13.2 per 1,000 live births,  $p < 0.001$ ), and perinatal death rate (21.2 vs. 13.2 per 1,000 total births,  $p < 0.001$ ) were statistically significantly higher among Syrian births compared to Jordanian births. The distribution of stillbirths, neonatal deaths and perinatal deaths for babies born to Syrian women according to gestational age and birth weight are shown in

**Figure 3.** The vast majority of deaths occurred among babies born prematurely or with low birth weight.

### Multivariate Analysis

In the bivariate analysis, the proportion of babies born with low birthweight or SGA, stillbirth rate, neonatal death rate, and perinatal death rate was statistically significantly higher among babies born to Syrian women than those born to Jordanian women. After adjusting for sociodemographic and obstetric characteristics of women (**Table 3**), only perinatal death was



**TABLE 3 |** Multivariate analysis of differences in birth outcomes for Jordanian newborns versus Syrian newborns (Jordan, May 2019 and December 2020).

	Univariate analysis			Multivariate analysis*				
	OR	95% confidence interval	p-value	OR	95% confidence interval	p-value		
Preterm birth	1.1	1.0	1.2	0.082	1.0	0.9	1.2	0.936
Low birthweight	1.2	1.1	1.4	<0.001	1.1	0.9	1.2	0.243
Small for GA	1.2	1.1	1.4	<0.001	1.1	1.0	1.3	0.081
Stillbirth	1.5	1.2	2.1	0.004	1.3	0.9	1.9	0.181
Neonatal death	1.6	1.3	2.1	<0.001	1.2	0.9	1.7	0.271
Perinatal death	1.6	1.3	2.0	<0.001	1.3	1.1	1.7	0.035

\*Adjusted for mother’s age, employment status, education, income, baby gender, multiplicity, parity, history of neonatal death, and history of stillbirth.

statistically significantly higher among Syrian babies compared to Jordanian babies (OR = 1.3, 95% CI: 1.1–1.7, *p* = 0.035).

## DISCUSSION

This study is the first to systematically assess and compare wide-ranging birth outcomes and disparities among Syrian women in the host community of Jordan. Using comprehensive multicentre data, our analysis specifically provided information about the discrepancies in LBW (<2,500 g), SGA, preterm delivery, birthweight, stillbirth, and perinatal death rate among Syrian and Jordanian women’s births.

Our findings indicate remarkable disparities in pregnancy outcomes among the Syrian women refugees in the host communities of Jordan. Compared to the Jordanian women, the adverse pregnancy outcomes were remarkably higher among the Syrian women refugees, a gravely concerning finding that resonated with a previous retrospective study conducted at two governmental hospitals in northern Jordan [24]. The previous study documented a significant inequality in terms of the rate of caesarean section, anaemia, and lower neonates’ APGAR scores and birthweight among Syrian women deliveries compared to those from Jordan [24]. When summed together, these findings indicate a priority for

more effective and tailored maternal-newborns healthcare services directed toward the refugee mothers in Jordan. Transformation is required in the current healthcare policies and interventions implemented in Jordan to improve women refugee’s access to and quality of antenatal, birth, and postnatal healthcare services.

Neonatal health outcomes, in general, have improved considerably in Jordan, as indicated by a slow but progressive decline in neonatal death rates over the past 20 years [25]. According to international reports, Jordan’s overall neonatal mortality rate is recorded at 9 per 1,000 LB [25], and above 14 per 1,000 LB, according to research-based information conducted nationally [19, 26]. Comparing our findings to these national reports, the neonatal and perinatal death rates recorded among Syrian births are remarkably higher than those reported nationally and internationally [19, 25, 26]. Adverse pregnancy outcomes are significantly high among Syrian women refugees on all measured parameters in this study. Our analysis showed that the overall birth outcomes among the Syrian women refugees in Jordan are poor and create a threat to achieving Jordan’s national and international development sustainability goals [27, 28].

Our findings regarding certain birth outcomes (e.g., weight at birth, SGA, low birthweight) were in agreement with those that have been reported among displaced Syrian women in a few

comparable studies from Lebanon [6, 29] and Turkey [2, 5, 30]. However, in opposition to our findings, numerous studies from Turkey demonstrated no disparities affecting Syrian women concerning caesarean birth rates [2, 30–33], preterm delivery [30, 32, 33], stillbirth [5, 34], abortion [32, 33], LBW (<2,500 g) [33, 35], and small for gestational birthweight [5]. These discrepancies could suggest that health policies and interventions that facilitate women refugees' access to and the provision of quality maternal and neonatal services could positively reduce the prevalence and disparities of adverse pregnancy outcomes among women refugees. An international conclusion should be drawn with caution and should be interpreted within its terrestrial context, considering the variations in the quality of healthcare services across countries.

Our findings also showed that other disparities also existed in the sociodemographic characteristics that are related to birth outcomes of the mothers from Syria compared to those from Jordan, specifically in terms of younger age, income, unemployment, and educational attainment. The results showed that the difference can be attributed to the background variables among the Jordanian and Syrian women. After controlling for these factors, the difference between the two groups of women seems to become not significant. These sociodemographic discrepancies could have contributed to worsening pregnancy outcomes and widening the healthcare disparity gap among the Syrian women refugees in Jordan. Low socioeconomic state [6], early-age/adolescent pregnancy [35], infrequent antenatal visits [6, 32, 34, 35], and the late arrival to skilled maternal care at birth [5, 24, 35] are contributing factors to adverse pregnancy outcomes that were prevailing among women refugees from Syria in other countries. The adverse birth outcomes are multifactorial, and this should be considered when designing and delivering care services to women refugees in the host countries.

Adolescent pregnancy, in particular, has been consistently observed to be prevalent among Syrian displaced women compared to native women in host countries [2, 35–38], which is in accordance with our comparison in Jordan. Adolescent pregnancy is one crucial factor associated with adverse birth outcomes worldwide [39, 40], which is similar to the case of Jordan [41]. The socioeconomic and healthcare needs, risks and pregnancy outcomes of Syrian adolescents and young women should receive further attention through humanitarian field assessments and research work. Our findings sheds light on the importance of designing healthcare policies and services to provide additional support for the vulnerable groups of women refugees.

## REFERENCES

- Giovetti O. *Forced Migration: 6 Causes and Examples* (2021). Available From: <https://www.concernusa.org/story/forced-migration-causes/> (Accessed August 12, 2021).
- Çelik İH, Arslan Z, Ulubaş Işık D, Tapısız ÖL, Mollamahmutoglu L, Baş AY, et al. Neonatal Outcomes in Syrian and Other Refugees Treated in a Tertiary Hospital in Turkey. *Turkish J Med Sci* (2019) 49(3):815–20. doi:10.3906/sag-1806-86

## Strengths and Limitations

Retrospective studies on exposing data are efficient means of gathering data in public health studies; however, researchers are limited with the nature and amount of the data excited. Although the current analysis was conducted in five leading referral hospitals in Jordan, the findings might only be transferred to some Jordanian women and Syrian refugees that might receive care outside these hospitals in the country. Despite this limitation, this study is the first multicentre study in Jordan to compare wide-ranging sociodemographic and birth outcomes between displaced Syrian refugees and Jordanian women. This study pave the way for future improvement in obstetric, neonatal, and birth outcomes and there related surveillance system.

## Conclusion

Syrian refugee mothers had higher rates of adverse birth outcomes and inequalities than women in Jordan's host communities. Ongoing evaluation of pregnancy outcomes is essential to enhance our understanding of the healthcare disparities among refugees in the host communities. However, it is more important to foster a comprehensive understanding of the multilevel contributing factors to help develop strategies on micro and macro levels for controlling these factors. Our findings suggest a healthcare priority for better improvement in the policies and provision of maternal-neonatal care services to women refugees in Jordan. Healthcare policies and services may be designed to provide additional support for these vulnerable groups of women refugees.

## ETHICS STATEMENT

The studies involving human participants were reviewed and approved by the Jordan University of Science and Technology. Written informed consent for participation was not required for this study in accordance with the national legislation and the institutional requirements.

## AUTHOR CONTRIBUTIONS

All authors listed have made a substantial, direct, and intellectual contribution to the work and approved it for publication.

## CONFLICT OF INTEREST

The authors declare that they do not have any conflicts of interest.

- Harakow HI, Hvidman L, Wejse C, Eiset AH. Pregnancy Complications Among Refugee Women: A Systematic Review. *Acta Obstetrica Gynecologica Scand* (2021) 100(4):649–57. doi:10.1111/aogs.14070
- Gibson-Helm ME, Teede HJ, Cheng IH, Block AA, Knight M, East CE, et al. Maternal Health and Pregnancy Outcomes Comparing Migrant Women Born in Humanitarian and Nonhumanitarian Source Countries: A Retrospective, Observational Study. *Birth* (2015) 42(2):116–24. doi:10.1111/birt.12159
- Ozel S, Yaman S, Kansu-Celik H, Hancerliogullari N, Balci N, Engin-Ustun Y. Obstetric Outcomes Among Syrian Refugees: A Comparative Study at a

- Tertiary Care Maternity Hospital in Turkey. *Rev Bras Ginecol Obstet* (2018) 40(11):673–9. doi:10.1055/s-0038-1673427
6. Elshal H, Chour M, Halim SA, Kharpoutli S, Raad H. Factors Affecting Pregnancy Outcome in Refugee Mothers in Lebanon. *BAU J - Health Well-Being* (2021) 3(2):1–7. doi:10.54729/2789-8288.1128
  7. World Vision. *Syrian Refugee Crisis: Facts, FAQs, and How to Help* WA. USA: World Vision, Inc (2021).
  8. Regional Refugee & Resilience Plan. *Regional Strategic Overview 2021-2022: RSO* (2021). Available From: <http://www.3rpsyriacrisis.org/wp-content/uploads/2020/12/RSO2021.pdf> (Accessed July 24, 2021).
  9. Operational Data Portal. *Refugee Situation. Syria Regional Refugee Response -Jordan: Operational Data Portal 2021* (2021). Available From: <https://data2.unhcr.org/en/situations/syria/location/36> (Accessed July 24, 2021).
  10. Brown H, Giordano N, Maughan C, Wadeson A. *The 2019 Vulnerability Assessment Framework (VAF). The UN Refugee Agency (UNHCR), Action Against Hunger UK's Monitoring, Evaluation and Learning Services*. Geneva, Switzerland: International Labour Organisation (2019).
  11. Grameen Cr dit Agricole Foundation (GCAF), Swedish International Development Cooperation Agency (SIDA), United Nations High Commissioner for Refugees (UNHCR). *Assessing the Needs of Refugees for Financial and Non-Financial Services—Jordan* (2018). Available From: <https://www.unhcr.org/fr-fr/en/media/assessing-needs-refugees-financial-and-non-financial-services-jordan> (Accessed July 24, 2021).
  12. UNHCR. *Health-Refugees From All Nationalities: UNHCR; 2021* (2021). Available From: <https://help.unhcr.org/jordan/en/helpful-services-unhcr/health-services-unhcr/> (Accessed July 24, 2021).
  13. Al-Rousan T, Schwabkey Z, Jirmanus L, Nelson BD. Health Needs and Priorities of Syrian Refugees in Camps and Urban Settings in Jordan: Perspectives of Refugees and Health Care Providers. *East Mediterr Health J* (2018) 24(3):243–53. doi:10.26719/2018.24.3.243
  14. Amiri M, El-Mowafi IM, Chahien T, Yousef H, Kobeissi LH. An Overview of the Sexual and Reproductive Health Status and Service Delivery Among Syrian Refugees in Jordan, Nine Years Since the Crisis: A Systematic Literature Review. *Reprod Health* (2020) 17(1):166–20. doi:10.1186/s12978-020-01005-7
  15. Clark CJ, Spencer RA, Khalaf IA, Gilbert L, El-Bassel N, Silverman JG, et al. The Influence of Family Violence and Child Marriage on Unmet Need for Family Planning in Jordan. *J Fam Plann Reprod Health Care* (2017) 43(2):105–12. doi:10.1136/fjprhc-2014-101122
  16. Khader YS, Alyahya M, El-Khatib Z, Batieha A, Al-Sheyab N, Shattnawi K. The Jordan Stillbirth and Neonatal Mortality Surveillance (JSANDS) System: Evaluation Study. *J Med Internet Res* (2021) 23(7):e29143. doi:10.2196/29143
  17. Khader YS, Shattnawi KK, Al-Sheyab N, Alyahya M, Batieha A. The Usability of Jordan Stillbirths and Neonatal Deaths Surveillance (JSANDS) System: Results of Focus Group Discussions. *Arch Public Health* (2021) 79(1):29–10. doi:10.1186/s13690-021-00551-1
  18. Khader YS, Alyahya M, Batieha A, Tawel A. JSANDS: A Stillbirth and Neonatal Deaths Surveillance System. In: 2019 IEEE/ACS 16th International Conference on Computer Systems and Applications (AICCSA); Nov. 3 2019 to Nov. 7 2019; Abu Dhabi, United Arab Emirates (2019).
  19. Al-Sheyab NA, Khader YS, Shattnawi KK, Alyahya MS, Batieha A. Rate, Risk Factors, and Causes of Neonatal Deaths in Jordan: Analysis of Data From Jordan Stillbirth and Neonatal Surveillance System (JSANDS). *Front Public Health* (2020) 8:595379. doi:10.3389/fpubh.2020.595379
  20. Shattnawi KK, Khader YS, Alyahya MS, Al-Sheyab N, Batieha A. Rate, Determinants, and Causes of Stillbirth in Jordan: Findings From the Jordan Stillbirth and Neonatal Deaths Surveillance (JSANDS) System. *BMC Pregnancy and Childbirth* (2020) 20(1):571–8. doi:10.1186/s12884-020-03267-2
  21. World Health Organization. *The International Statistical Classification of Diseases and Health Related Problems ICD-10: Tenth Revision*. Geneva, Switzerland: World Health Organization (2004).
  22. Howson CP, Kinney MV, McDougall L, Lawn JE. Born Too Soon: Preterm Birth Matters. *Reprod Health* (2013) 10(1):1–9. doi:10.1186/1742-4755-10-S1-S1
  23. World Health Organization. *The WHO Application of ICD-10 to Deaths During the Perinatal Period: ICD-PM*. Geneva, Switzerland: World Health Organization (2016).
  24. Alnuaimi K, Kassab M, Ali R, Mohammad K, Shattnawi K. Pregnancy Outcomes Among Syrian Refugee and Jordanian Women: A Comparative Study. *Int Nurs Rev* (2017) 64(4):584–92. doi:10.1111/inr.12382
  25. UNICEF. *Country Profiles: Jordan 2021* (2021). Available From: <https://data.unicef.org/country/jor/> (Accessed March 27, 2022).
  26. Batieha AM, Khader YS, Berdzuli N, Chua-Oon C, Badran EF, Al-sheyab NA, et al. Level, Causes and Risk Factors of Neonatal Mortality, in Jordan: Results of a National Prospective Study. *Matern Child Health J* (2016) 20(5):1061–71. doi:10.1007/s10995-015-1892-x
  27. World Health Organization. *Global Strategy for Women's, Children's and Adolescents' Health (2016–2030): early childhood development* (2018). Available From: [https://iris.who.int/bitstream/handle/10665/276423/A71\\_19Rev1-en.pdf?sequence=1](https://iris.who.int/bitstream/handle/10665/276423/A71_19Rev1-en.pdf?sequence=1) (Accessed July 24, 2021).
  28. United Nations. *United Nations Jordan; 2022* (2022). Available From: <https://jordan.un.org/en/sdgs/3> (Accessed July 24, 2021).
  29. Abdulrahim S, El Rafei R, Beydoun Z, El Hayek GY, Nakad P, Yunis K. A Test of the Epidemiological Paradox in a Context of Forced Migration: Low Birthweight Among Syrian Newborns in Lebanon. *Int J Epidemiol* (2019) 48(1):1024–86. doi:10.1093/ije/dyz087
  30. Vural T, G lbaşı C, Bayraktar B, G lbaşı H, Yıldırım AG. Are Syrian Refugees at High Risk for Adverse Pregnancy Outcomes? A Comparison Study in a Tertiary Center in Turkey. *J Obstet Gynaecol Res* (2021) 47(4):1353–61. doi:10.1111/jog.14673
  31. G ng r ES, Seval O, İlhan G, Verit FF. Do Syrian Refugees Have Increased Risk for Worse Pregnancy Outcomes? Results of a Tertiary Center in İstanbul. Suriyeli M ltecilerin Daha K t  Gebelik Sonu ları A ısından Artmıř Riskleri Var Mıdır? *İstanbul'da Tersiye bir Merkezin Sonu ları* (2018) 15(1):23–7. doi:10.4274/ijod.64022
  32. Kiyak H, Gezer S, Ozdemir C, Gunkaya S, Karacan T, Gedikbasi A. Comparison of Delivery Characteristics and Early Obstetric Outcomes Between Turkish Women and Syrian Refugee Pregnancies. *Niger J Clin Pract* (2020) 23(1):12–7. doi:10.4103/njcp.njcp\_10\_18
  33. Sayılı U, Ozgur C, Bulut Gazanfer O, Solmaz A. Comparison of Clinical Characteristics and Pregnancy and Neonatal Outcomes Between Turkish Citizens and Syrian Refugees With High-Risk Pregnancies. *J Immigrant Minor Health* (2021) 24:1177–85. doi:10.1007/s10903-021-01288-3
  34. Kanmaz AG, İnan AH, Beyan E,  zg r S, Budak A. Obstetric Outcomes of Syrian Refugees and Turkish Citizens. *Arch Iranian Med* (2019) 22(9):482–8.
  35. Erenel H, Aydoğan Mathyk B, Sal V, Ayhan I, Karatas S, Koc Bebek A. Clinical Characteristics and Pregnancy Outcomes of Syrian Refugees: A Case-Control Study in a Tertiary Care Hospital in İstanbul, Turkey. *Arch Gynecol Obstet* (2017) 295(1):45–50. doi:10.1007/s00404-016-4188-5
  36. Bacanaklı BH, Kaban I, Yildirim SG, Sinaci E. Birth Characteristics of Syrian Refugees at a Tertiary Medical Center in İstanbul, Turkey. *Eur J Obstet Gynecol Reprod Biol* (2019) 234:e140–e. doi:10.1016/j.ejogrb.2018.08.466
  37. Dopfer C, Vakılzadeh A, Happle C, Kleinert E, M ller F, Ernst D, et al. Pregnancy Related Health Care Needs in Refugees—A Current Three Center Experience in Europe. *Int J Environ Res Public Health* (2018) 15(9):1934. doi:10.3390/ijerph15091934
  38. Turkyay  , Aydin  ,  alıřkan E, Salıcı M, Terzi H, Astepe B. Comparison of the Pregnancy Results Between Adolescent Syrian Refugees and Local Adolescent Turkish Citizens Who Gave Birth in Our Clinic. *J Maternal-Fetal Neonatal Med* (2020) 33(8):1353–8. doi:10.1080/14767058.2018.1519016
  39. Wong SP, Twynstra J, Gilliland JA, Cook JL, Seabrook JA. Risk Factors and Birth Outcomes Associated With Teenage Pregnancy: A Canadian Sample. *J Pediatr Adolesc Gynecol* (2020) 33(2):153–9. doi:10.1016/j.jpag.2019.10.006
  40. Karatařlı V, Kanmaz AG, İnan AH, Budak A, Beyan E. Maternal and Neonatal Outcomes of Adolescent Pregnancy. *J Gynecol Obstet Hum Reprod* (2019) 48(5):347–50. doi:10.1016/j.jogoh.2019.02.011
  41. Khader YS, Batieha A, Al-Hader R, Sa'ad SH, Hijazi SS. Rate of Teenage Pregnancy in Jordan and Its Impact on Maternal and Neonatal Outcomes. *Int J Adolesc Med Health* (2019) 31(6). doi:10.1515/ijamh-2017-0075

Copyright   2023 Al-Shatanawi, Khader, Abdel Razeq, Khader, Alfaqih, Alkouri and Alyahya. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.